### **UNCLASSIFIED**

# AD NUMBER AD467863 **NEW LIMITATION CHANGE** TO Approved for public release, distribution unlimited **FROM** Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; Dec 1964. Other requests shall be referred to Office of Naval Research, Arlington, VA. **AUTHORITY** ONR ltr, 13 Sep 1977

THIS REPORT HAS BEEN DELIMITED

AND CLEARED FOR PUBLIC RELEASE

UNDER DOD DIRECTIVE 5200.20 AND

NO RESTRICTIONS ARE IMPOSED UPON

ITS USE AND DISCLOSURE.

DISTRIBUTION STATEMENT A

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

## SECURITY MARKING

The classified or limited status of this report applies to each page, unless otherwise marked.

Separate page printouts MUST be marked accordingly.

THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE OF THE UNITED STATES WITHIN THE MEANING OF THE ESPIONAGE LAWS, TITLE 18, U.S.C., SECTIONS 793 AND 794. THE TRANSMISSION OR THE REVELATION OF ITS CONTENTS IN ANY MANNER TO AN UNAUTHORIZED PERSON IS PROHIBITED BY LAW.

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

CATALOGED BY: DEN 67863

# FROM HIGH PRESSURE GASES

by

Ralph Zirkind





December 1964

### POLYTECHNIC INSTITUTE OF BROOKLYN

DEPARTMENT

of
AEROSPACE ENGINEERING

and
APPLIED MECHANICS

PIBAL REPORT NO. 814

# EMISSION OF OPTICAL RADIATION FROM HIGH PRESSURE GASES

by

Ralph Zirkind

This research was conducted under the sponsorship of the Office of Naval Research under Contract No. Nonr 839(34), Project No. NR 061-135.

Reproduction in whole or in part is permitted for any purpose of the United States Government.

Polytechnic Institute of Brooklyn

Department

of

Aerospace Engineering and Applied Mechanics

December 1964

PIBAL Report No. 814

### EMISSION OF OPTICAL RADIATION FROM HIGH PRESSURE GASES†

by

Ralph Zirkind\*
Polytechnic Institute of Brooklyn

#### SUMMARY

The observed spectral emittance of high temperature and density nitrogen was examined where the gas was compressed isentropically to a temperature and pressure of  $7200^{\circ}$ K and  $4500 \text{ kg/cm}^2$  respectively. The analysis indicated that the radiation was composed of a near black body continuum and of nitrogen band emissions where the predominant bands were  $N_2^+(1-)$ ,  $N_2(1+)$  and  $N_2(2+)$ . Examination of the ratio of the latter two bands then made possible the reconstruction of the temperature history of the gas in the vicinity of peak compression. It was determined also that in comparison to the temperature, the relatively high density ratio of 250 played a minor role in the emitted radiant intensity.

This research was conducted under the sponsorship of the Office of Naval Research under Contract No. Nonr 839(34), Project No. NR 061-135.

\*Visiting Professor, Aerospace Engineering

#### SECTION I

#### INTRODUCTION

The ability to compress a gas, isentropically or adiabatically, permits the examination of the role that density plays in the spectral emittance of the compressed gas at optical frequencies.

In reference 1, Vodar and Vu reviewed the recent literature which pertains to the effects of pressure upon spectral line and band broadening; however, no considerations were included relative to density effects. In references 2 and 3, research on gases isentropically compressed to pressures as high as 8000 kg/cm<sup>2</sup> were reported. The methods used by both investigators are similar; that is, a piston is driven ballistically into a closed cylinder. The relationships of the state variables for fixed pressure ratio functions of  $\gamma$ , the ratio of specific heats, are

$$\frac{T_1}{T_0} = \frac{P_1}{P_0} \qquad \text{and} \qquad \frac{\rho_1}{\rho_0} = \frac{P_1}{P_0} \frac{1/\gamma}{\rho_0}$$
 (1)

where P,  $\rho$  and T have their usual meaning and subscripts "l" and "o" refer to final and initial conditions respectively. Now the value of  $\gamma$  is greatest for a monatomic gas, 5/3, and therefore can produce the highest temperature. In both instances argon gas was the working fluid.

The effects of adiabatic compression will be studied through the results obtained from the utilization of shock tubes. The real difference between the methods of compression is that for the same pressure ratio, the Rankine-Hugoniot conditions preclude a density ratio greater than ~ 4;

The work in the United States at NOL, White Oaks, is relatively new in comparison to that in the U.S.S.R.<sup>2</sup> and apparently is conducted without knowledge of it.

whereas, isentropically, a value of 250 can be achieved. That is, under the assumption of no ionization in argon (valid for shock velocities  $M_1 < 10$ ) the density ratio is given by Eq. (2),

$$\frac{\rho_1}{\rho_0} = \frac{(\gamma + 1)M_1^2}{(\gamma - 1)M_1^2 + 2} . \tag{2}$$

Hence, for  $\gamma = 1.6$ . the asymptotic value for  $\rho_1/\rho_0$  is approximately 4, which is significantly different from the value in the isentropic case.

Since the intensity of absorption varies as the square of the density, one might expect a notable difference between the two cases.

The paper will present the experimental data and an analysis the reof, followed by a discussion of the results and comparison with theory.

## SECTION II EXPERIMENTAL DATA

In reference 2 the results of an experiment on high pressure industrial argon were reported. The constituents of the analyzed gas were found to be as follows: Argon - 94%:  $N_2$  - 4%;  $H_2O$  - 2%, and trace amounts of  $O_2$ . A maximum compression of 4500 kg/cm<sup>2</sup> was achieved in the experiment discussed here and the duration of radiation emission was 200 microseconds. The thermodynamic relation between pressure and temperature indicates that the gas temperature at peak compression is about 7200 K.

The published data in reference 2 are not complete; however, a sufficient quantity are available for meaningful analysis.

In Fig. 1, the intensity is plotted as a function of wavelength in the spectral range of  $\lambda\lambda$  265.0 - 420.0  $m_{tl}$  at three specific times, maximum compression and  $\neq$  10 microseconds thereof. The second set of data are time-resolved spectra which yielded an indication of the available species and contaminants, e.g., OH. The time-resolved spectrograms obtained prior to maximum compression, namely, those at 45 and 52 microseconds and the one at maximum compression, 100 microseconds, provided a

measure of  $N_2(2+)$  and  $N_2(1+)$ .

The intensity ratio of the 428.0 mu and 568.0 mu bands of  $N_2$  (2+) and  $N_3$  (1+), as estimated from reference 2, is given in Table 1. The estimate provides for the contribution due to the continuum. In Fig. 2, the data of Table 1 are plotted together with calculated results.

The first and second positive  $N_2$  bands were selected for study since they were readily identified and sufficiently displaced in wavelength.

TABLE 1

Time Dependence of the Intensity Ratio of N<sub>2</sub>(1+)/N<sub>2</sub>(2+)

Time (microseconds)	1(5680)/1(4280)
48	5.00
52	3.33
100	2.0

### SECTION III DISCUSSION

In Fig. 1, the relative spectral intensity at several wavelengths, 262, 362, and 410 mu, for several times, maximum compression  $\neq$  10 usecs are compared to the spectral intensities expected from a black body at 5500-7200°K. The latter value corresponds to the temperature at maximum compression. In the spectral region of 2600 mu to 3600 mu the slope of both the pre-maximum and post-maximum data points, when normalized to the intensity at 262 mu, yields a temperature of about 6500°K. At maximum compression, the slope would correspond to approximately 7200°K. The real departure from a black body occurs for the wavelength value of 4100 mu. The possible contributions at this wavelength are (1)  $N_s^+$ (1-) and (2) NO B. The latter possibility may be discarded for the following two reasons:

- (1) the spectrograms show a marked intensity drop between 470-540 mu, which would be filled in if NO was significant, and
- (2) above 4000°K, the temperature regime of interest here, NO commences to dissociate, the dissociation becoming nearly complete at ~6500°K.

If the contribution of  $N_2^+(l-)$  is subtracted from the total, the value at 410  $m_U$  drops to ~19 and the data points behave more nearly like a black body.

In Fig. 2, the calculated relative intensity ratio of  $N_2$  (1+) to  $N_2$  (2+) at wavelengths 568, 428 mu respectively, are plotted as a function of temperature and density ratio. The experimental values, Table 1, obtained at times equal to maximum compression ( $\rho/\rho_0 \sim 250$ ) and somewhat earlier times, are also presented. The value at maximum compression gives a temperature value of  $7250^{\circ}$  K, which corresponds very closely to the expected temperature of the compressed Argon gas. The temperature of the gas, as estimated from Fig. 2, would be about  $6000^{\circ}$  K and  $5100^{\circ}$  K, which is reasonable. These data further support the argument made above, that NO contributes little or no radiation to the total observed radiation.

# SECTION IV CONCLUDING REMARKS

The observed optical radiation from industrial argon (argon plus nitrogen) has been examined and analyzed. From a comparison of the time-resolved spectra, the relative contributions from the continuum and band emitters have been estimated.

The first result obtained is that the compressed argon gas behaves like a black body emitter where the temperatures correspond to those values calculated from the standard laws for ideal gases. Secondly, NO-8 does not contribute to the radiation as expected, and the primary band emitters are  $N_2(2+)$  and  $N_2(1+)$  with a contribution from  $N_2(1-)$ . Finally,

the optical behavior of gas at densities ratio as high as 250 is still primarily determined by the gas temperature.

The experimental method discussed in this paper would provide an interesting procedure for the study of electronic excitation of molecules. The latter area is an important region and at present inadequately covered.

#### SECTION V

#### REFERENCES

- 1. Vodar, B. and Vu, H.: <u>Intensites Absolues des Transitions Incluites par La Pression</u>. J. Quant. Spectro. Radiative Transfer, 3, 397, 1963.
- 2. Ryabinin, Y.N.: Gases at High Densities and Temperatures.
  Pergamon Press, 1961.
- 3. Lalos, G. T. and Hammond, G. L.: Emission Spectra of Hot Dense Gases. Astrophysical J., 135, 616, 1962.





FIG. I. RELATIVE INTENSITY VS & FOR SEVERAL TIMES



これのあれているのであることはないのできますのできますのできます。

FIG. 2. BAND INTENSITY RATIO AS A FUNCTION OF TEMPERATURE

Unclassified
Security Classification

Security Classification	· · · · · · · · · · · · · · · · · · ·		<u> </u>			
DOCUMENT C (Security classification of title, body of abstract and inde	ONTROL DATA - RE		the overall report i	e classified)		
1. ORIGINATING ACTIVITY (Corporate author) Polytechnic Institute of Brooklyn Graduate Center		24. REPORT SECURITY CLASSIFICATION				
Dept. of Aerospace Engineering & App. Route 110, Farmingdale, New York 117	lied Mechanics 35	25 GROU	P			
3. REPORT TITLE	•					
EMISSION OF OPTICAL RADIATION FROM P	HIGH PRESSURE GA	SES				
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)						
Interim report						
S. AUTHOR(S) (Leet name, first name, initial)						
Zirkind, Ralph		•.				
6. REPORT DATE	74. TOTAL NO. OF	AGES	75. NO. OF REP	78		
December 1964	7		3			
Se. CONTRACT OR SEANT NO. NOnz 839(34)	94 ORIGINATOR'S R	EPORT NUM	in ER(S)			
& PROJECT NO. 061-135	• PIBAL Repo					
				•		
<b>c.</b>	9b. OTHER REPORT	NO(8) (Any	other numbers that	may be sesigned		
10. A VAILABILITY/LIMITATION NOTICES				<del>.</del>		
Qualified requesters may obtain copic	es of this renor	t from 1	DDC .			
- Quartified requesters may obtain copie	es of this repor	· IIOm				
11. SUPPLEMENTARY NOTES	12. SPONSORING MILI	TARY ACT	VITY			
S S S S S S S S S S S S S S S S S S S	Office of Naval Research					
	Department of the Navy					
	Washington	, D. C.	20360			

13. ABSTRACT

The observed spectral emittance of high temperature and density nitrogen was examined where the gas was compressed isentropically to a temperature and pressure of  $7200^{\circ}$ K and  $4500 \text{ kg/cm}^2$  respectively. The analysis indicated that the radiation was composed of a near black body continuum and of nitrogen band emissions where the predominant bands were N2<sup>†</sup>(1-), N2(1+) and N2(2+). Examination of the ratio of the latter two bands then made possible the reconstruction of the temperature history of the gas in the vicinity of peak compression. It was determined also that in comparison to the temperature, the relatively high density ratio of 250 played a minor role in the emitted radiant intensity.

DD 15084 1473

Unclassified
Security Classification

	_								
C.				~	מו		fine	tion	
.31	76.	411	LV	٠.		881		i Li Un	

14.		LINK A		LINK B		LINK E		
KEY WORDS		, no	DLK	WT	ROLE	wT	ROLE	WT
Optical Radiation			1					: 
Gases	•							
			Ì					
		i						
			Ì	Ì				
			İ					
			1	İ				
		ŀ			;		į	
		l						
								}

#### INSTRUCTIONS

- 1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (comporate author) issuing the report.
- 24. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.
- 2b. GROUP: Automatic downgrading is specified in DoD Directive 5200, 10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authoriked.
- 3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.
- DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.
- 5. AUTHOR(5): If ter the name(s) of author(s) as shown on or in the report. Enter isst name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement
- REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.
- 7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 76. NUMBER OF REFERENCES: Enter the total number of references cited in the report.
- So. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.
- Sh. Sc. & Sd. PROJECT NUMBER: Enter the appropriate military department identification, such as project number. subproject number, system numbers, task number, etc.
- 9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (either by the originator or by the aponeor), also enter this number(s).
- AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorised."
- "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through
- (5) "All distribution of this report is controlled Qualified DDC users shall request through

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

- IL SUPPLEMENTARY NOTES: Use for additional explanatory notes.
- 12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.
- 13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical re-port. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (73). (5), (C), or (U).

There is no limitation on the length of the shatract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rales, and weights is optional.